



Candida albicans Impacts *Staphylococcus aureus* Alpha-Toxin Production via Extracellular Alkalinization

Olivia A. Todd,^a Mairi C. Noverr,^b Brian M. Peters^{c,d}

^aIntegrated Program in Biomedical Sciences, Microbiology, Immunology, and Biochemistry track, University of Tennessee Health Science Center, Memphis, Tennessee, USA

^bDepartment of Microbiology and Immunology, Tulane University, New Orleans, Louisiana, USA

^cDepartment of Clinical Pharmacy and Translational Science, University of Tennessee Health Science Center, Memphis, Tennessee, USA

^dDepartment of Microbiology, Immunology, & Biochemistry, University of Tennessee Health Science Center, Memphis, Tennessee, USA

ABSTRACT *Candida albicans* and *Staphylococcus aureus* are common causes of nosocomial infections with severe morbidity and mortality. Murine polymicrobial intra-abdominal infection (IAI) with *C. albicans* and *S. aureus* results in acute mortality dependent on the secreted cytolytic effector alpha-toxin. Here, we confirmed that alpha-toxin is elevated during polymicrobial growth compared to monomicrobial growth *in vitro*. Therefore, this study sought to unravel the mechanism by which *C. albicans* drives enhanced staphylococcal alpha-toxin production. Using a combination of functional and genetic approaches, we determined that an intact *agr* quorum sensing regulon is necessary for enhanced alpha-toxin production during coculture and that a secreted candidal factor likely is not implicated in elevating *agr* activation. As the *agr* system is pH sensitive, we observed that *C. albicans* raises the pH during polymicrobial growth and that this correlates with increased *agr* activity and alpha-toxin production. Modulation of the pH could predictably attenuate or activate *agr* activity during coculture. By using a *C. albicans* mutant deficient in alkalinization (*stp2Δ/Δ*), we confirmed that modulation of the extracellular pH by *C. albicans* can drive *agr* expression and toxin production. Additionally, the use of various *Candida* species (*C. glabrata*, *C. dubliniensis*, *C. tropicalis*, *C. parapsilosis*, and *C. krusei*) demonstrated that those capable of raising the extracellular pH correlated with elevated *agr* activity and alpha-toxin production during coculture. Overall, we demonstrate that alkalinization of the extracellular pH by the *Candida* species leads to sustained activation of the staphylococcal *agr* system.

IMPORTANCE *Candida albicans* and *Staphylococcus aureus* are commonly coisolated from central venous catheters and deep-seated infections, including intra-abdominal sepsis. Thus, they represent a significant cause of nosocomial morbidity and mortality. Yet how these organisms behave in the context of polymicrobial growth remains poorly understood. In this work, we set out to determine the mechanism by which activation of the staphylococcal *agr* quorum sensing system and production of its major virulence effector alpha-toxin is enhanced during coculture with *C. albicans*. Surprisingly, we likely ruled out that a secreted candidal factor drives this process. Instead, we demonstrated that alkalinization of the extracellular milieu by *C. albicans* and other *Candida* species correlated with elevated *agr* activity. Thus, we propose a mechanism where modulation of the extracellular pH by fungal opportunists can indirectly alter virulence of a bacterial pathogen. Uncovering molecular events that drive interkingdom pathogenicity mechanisms may enhance surveillance and treatment for devastating polymicrobial infections.

KEYWORDS *Candida*, *Staphylococcus*, toxin, pH, polymicrobial, coculture

Citation Todd OA, Noverr MC, Peters BM. 2019. *Candida albicans* impacts *Staphylococcus aureus* alpha-toxin production via extracellular alkalinization. mSphere 4:e00780-19. <https://doi.org/10.1128/mSphere.00780-19>.

Editor Aaron P. Mitchell, Carnegie Mellon University

Copyright © 2019 Todd et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Brian M. Peters, brian.peters@uthsc.edu.

Received 25 October 2019

Accepted 1 November 2019

Published 13 November 2019

Candida albicans, an opportunistic polymorphic fungus, and *Staphylococcus aureus*, a ubiquitous bacterial pathogen, rank among the top organisms responsible for life-threatening invasive disease. Not only do these pathogens cause significant morbidity and mortality on their own, evidence for their existence as part of polymicrobial consortia has surfaced. For instance, *C. albicans* and *S. aureus* have been coisolated from a variety of biotic and abiotic surfaces, including central venous catheters, prosthetic implants, skin, and mucosal layers (1, 2). Moreover, they have been coassociated with several polymicrobial diseases, including burn wound superinfection, ventilator-associated pneumonia, urinary tract infection, cystic fibrosis, and bloodstream infection (3–6). However, perhaps the most comprehensive line of investigation of this particular coinfection has focused on their role during intra-abdominal infection (IAI).

IAI is a spectrum of diseases characterized by microbial invasion and subsequent inflammation of the abdominal cavity (7). Mortality rates for such infections typically range between 10 and 30%; however, mortality rates involving a fungal pathogen (e.g., *C. albicans*) can approach 80%, even with appropriate treatment (8, 9). Using a murine model of IAI, a series of studies by Carlson demonstrated that *C. albicans* enhanced the virulence of *S. aureus*, as coinfection reached 100% mortality within days postinfection (p.i.) while monomicrobial infection with either pathogen was nonlethal (10). Studies designed to titrate various inoculating doses of *C. albicans* and *S. aureus* during coinfection revealed that this apparent synergism was not mutual, as *C. albicans* virulence was not augmented by low doses of *S. aureus* (11). Synergistic mortality rates were found to be dependent on various toxins produced by *S. aureus* (12). Although these studies failed to delineate which specific *S. aureus* toxin was responsible, they established their important role in driving pathogenicity in the context of polymicrobial IAI.

Toxin expression in *S. aureus* is governed by a complex set of transcriptional regulators that respond to endogenous and environmental stimuli. Perhaps the best well-characterized mechanism is driven by the accessory gene regulator (*agr*) quorum sensing system that is activated in a cell density-dependent manner. The *agr* operon is composed of four genes encoding AgrA, AgrB, AgrC, and AgrD proteins (13). AgrD serves as the immature signal peptide that is proteolytically processed and secreted by membrane-bound AgrB, releasing the mature signal molecule autoinducing peptide 2 (AIP2). AIP2 can be sensed by the surface-bound receptor and histidine kinase AgrC that phosphorylates and activates the AgrA transcription factor. Activation of AgrA simultaneously upregulates expression of the *agr* operon completing a positive-feedback loop, while also ultimately downregulating colonization factors (e.g., adhesins) and upregulating virulence factors, including toxins (13, 14). While *agr* is undoubtedly induced as a consequence of quorum development, it is also highly susceptible to environmental factors, including high salt, glucose, subinhibitory antibiotic concentrations, and pH (15, 16).

Our laboratory has recently demonstrated that during *in vitro* growth, *C. albicans* has the capacity to activate the *agr* regulon, leading to exacerbated production of alpha-toxin, a potent staphylococcal virulence determinant capable of lysing a variety of host cells, causing tight-junction loss, and activating numerous innate proinflammatory pathways (17). Moreover, this virulence factor was crucial for driving lethal synergism during polymicrobial IAI. By using a combination of genetic and functional assays, the objective of this study was to attempt to elucidate the mechanism by which *C. albicans* activates *agr* signaling and alpha-toxin production by *S. aureus* and to determine the extent that other non-*albicans* *Candida* (NAC) species can augment alpha-toxin release (17). In the course of conducting these studies, we discovered that modulation of the extracellular pH by *C. albicans* creates an optimal environment for robust induction of the staphylococcal *agr* system, further highlighting how complex ecological signals may intersect with virulence during this prevalent fungal-bacterial interaction.




FIG 1 *C. albicans* enhances *S. aureus* alpha-toxin production. (A) Hemolytic activity of monomicrobial (*C. albicans* SC5314 [CA] or *S. aureus* JE2 [SA]) and polymicrobial cultures (CA plus SA [CA+SA]) was functionally assessed by depositing 20 μ l of cell-free culture supernatants into wells on a sheep blood agar plate. Images were taken after incubation at 37°C for 24 h using a digital scanner. (B) A P3-GFP reporter strain of *S. aureus* was incubated alone or with *C. albicans*. After 16 h of growth, 100 μ l culture was removed in triplicate and added to a 96-well plate, and fluorescence (in arbitrary fluorescence units [AFU]) was measured at 488/515 nm on a fluorimeter. (C) The concentration of alpha-toxin in supernatants from monomicrobial and polymicrobial cultures was measured by ELISA. Data are representative of three independent repeats and expressed as the means plus standard errors of the means (SEM) (error bars). Data were assessed for significance using one-way analysis of variance (ANOVA) and Dunnett's posttest. Values that are significantly different are indicated by asterisks as follows: **, $P < 0.01$; ***, $P < 0.001$.

RESULTS

***C. albicans* augments staphylococcal alpha-toxin production and does not complement *agr* signaling in *trans*.** Similar to previously reported findings (17), coculture of *C. albicans* and *S. aureus* led to elevated hemolytic toxin production compared to monoculture (Fig. 1A), as assessed by a functional hemolytic assay on sheep blood agar. Unsurprisingly, monoculture of *C. albicans* did not demonstrate lysis (Fig. 1A), as this fungus is not commonly reported to lyse red blood cells on microbiological agar. While the regulation of virulence factor production in *S. aureus* is multifactorial, the *agr* quorum sensing system plays a major role in governing increased toxin expression. In order to confirm that the *agr* system was more robustly activated during coculture, a P3-GFP (green fluorescent protein) reporter system (the P3 promoter is a target of phosphorylated and activated AgrA and ultimate driver of toxin expression) was employed. Indeed, the results of reporter analysis indicated an approximately 2.5-fold induction of *agr* (Fig. 1B), which correlated with ~4-fold production of alpha-toxin as measured by a specific enzyme-linked immunosorbent assay (ELISA) (Fig. 1C). These results were consistent with those in Fig. 1A, given that the hemolytic phenotype observed on sheep blood agar is dependent on alpha-toxin activity.

The next logical line of investigation was to determine whether *C. albicans* may be producing a protein or other small molecule that could be activating the *agr* system in *S. aureus*, leading to upregulation of toxin. Loss of *agrA* is predicted to largely ablate quorum sensing and elevated toxin production, as its activated form binds to both P2 and P3 promoters to drive the *agr* regulon and decrease repressor of toxin (*rot*), respectively. However, deletion of *agrB* would theoretically only attenuate secretion of the quorum signal peptide autoinducing peptide 2 (AIP2) (18–20). During monoculture, this would disrupt sensing of the quorum signal via AgrC and negatively impact toxin production. However, if *C. albicans* produced a molecule(s) that could be sensed by AgrC, then the native regulatory circuit could be bypassed during coculture, resulting in elevated toxin expression. Therefore, similar coculture assays were conducted as in Fig. 1, this time including isogenic Δ *agrA* and Δ *agrB* mutants. As predicted, disruption




FIG 2 *Candida*-enhanced toxin production is dependent on intact staphylococcal *agrA* and *agrB* signaling. (A) Hemolytic activity of polymicrobial cultures of *C. albicans* (CA) with wild-type *S. aureus* (SA) or isogenic strains lacking *agr* genes (Δ *agrA* or Δ *agrB*) was functionally assessed by depositing 20 μ l of cell-free culture supernatants into wells on a sheep blood agar plate. Images were taken after incubation at 37°C for 24 h using a digital scanner. (B) Levels of alpha-toxin were measured in polymicrobial culture supernatants by ELISA. Data are representative of three independent repeats and expressed as the means plus SEM. Data were assessed for significance using one-way ANOVA and Dunnett's posttest. Statistical significance: *, $P < 0.05$.

of *agrA* led to a nearly complete loss of hemolysis on sheep blood (Fig. 2A) and significantly reduced levels of alpha-toxin (Fig. 2B). Moreover, deletion of *agrB* demonstrated similar toxin phenotypes compared to the Δ *agrA* mutant (Fig. 2A and B). These results suggested that an intact *agr* regulon is required for elevated toxin expression. However, the augmented toxin phenotype was unlikely mediated by a fungal ligand-bacterial receptor interaction governed through AgrC sensing.

A role for extracellular pH in modulating *agr* signaling during coculture.

Previous reports have determined that regulation of *agr* is influenced by a number of physiologic factors, including low extracellular pH (15, 16). Therefore, we assessed the pH of monocultures and cocultures following standard growth conditions. The pH of *C. albicans* monoculture was estimated to be $\sim 7.5 \pm 0.1$, while that of *S. aureus* monoculture was $\sim 5.2 \pm 0.2$ (Fig. 3A). Interestingly, the pH of the coculture was $\sim 6.7 \pm 0.2$ (Fig. 3A). The pH of fresh Trypticase soy broth (TSB) plus 0.2% glucose (TSB-g) was determined to be 7.2 ± 0.2 (data not shown). Therefore, it appeared that *C. albicans* was elevating or maintaining the pH in a range which is optimal for *agr* activation. We next determined whether enhanced *agr* activity during coculture could be overridden by experimental modulation of the pH via buffering TSB-g with 100 mM morpholinepropanesulfonic acid (MOPS). We specifically chose to assess pH points that closely matched those representing *C. albicans* monoculture, *S. aureus* monoculture, and coculture values. Similar to previous results, use of the P3-GFP reporter system indicated induction of *agr* signaling in unbuffered medium (Fig. 3B). However, when the pH was buffered to acidic conditions (pH 5.5), *agr* signaling was significantly attenuated (Fig. 3B). When the pH was set to the *agr* optimum (pH 6.5), much higher levels of *agr* signaling were observed; however, differences between mono and coculture were no longer distinguishable (Fig. 3B). Similar results were found at a slightly more alkaline pH (pH 7.5), where *agr* signaling between monoculture and cocultures was indiscernible (Fig. 3B). These results suggest that extracellular pH strongly influences *agr* signaling *in vitro* and likely drives augmentation of toxin production during coculture.

In order to rule out the possibility that growth rates between different pH conditions were impacting *agr* reporter results, aliquots of each culture were taken at the experimental endpoint and plated onto selective microbiological media to enumerate both *C. albicans* and *S. aureus*. Although there was slight variation between each condition, there was no significant difference in the number of colonies of *C. albicans* (Fig. 3C) recovered during monoculture and coculture. Moreover, there was no significant difference between growth in buffered or unbuffered media. The same finding was true for *S. aureus* (Fig. 3C) during monoculture and coculture. Collectively, these results suggest that extracellular pH and not microbial growth accounts for disparate *agr* activity observed during mono- and coculture.

In order to rule out the possibility that a factor secreted by *C. albicans* activates *agr* signaling, *S. aureus*(pDB22) was grown at pH 5.5 overnight to attenuate *agr* signaling.




FIG 3 Synergistic hemolysis is partially dependent on extracellular pH and can be overridden by its modulation. (A) The pH of monomicrobial (SA or CA) and polymicrobial (CA+SA) cultures was measured after 16 h of incubation at 37°C. (B) Assays were conducted similarly as in panel A, except that the pH of TSB-g was buffered as indicated. Activation of staphylococcal *agr* was assessed using a GFP reporter assay. (C) Aliquots from each culture were taken at the endpoint to measure microbial counts by plating on selective media. Blue bars indicate *C. albicans* counts, and yellow bars indicate *S. aureus* counts. Solid color bars depict counts from monomicrobial (mono) cultures, while hatched bars depict counts from polymicrobial (poly) cultures. (D) Buffered (pH 7.0) spent CA and SA culture supernatants or fresh media were added to *S. aureus*(pDB22) initially grown at pH 5.5. Fluorescence (488 nm) was captured kinetically to assess *agr* activation. All data are representative of three independent repeats and expressed as the means ± SEM. Data were assessed for significance using one-way ANOVA and Dunnett’s posttest. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Spent culture supernatants from *C. albicans* or *S. aureus* JE2 grown at pH 7 were filter sterilized, reconstituted in fresh concentrated growth medium, and added to *S. aureus*-(pDB22) to elicit *agr* activation. Fresh culture medium was also used as a control. Results demonstrated that while spent culture medium from strain JE2 more rapidly and robustly activated the *agr* system, the addition of spent *C. albicans* supernatant did not give results different from those after the addition of fresh culture medium, suggesting that culture pH and not a specific candidal factor drives quorum signaling (Fig. 4D).

Several recent reports have demonstrated that *C. albicans* possesses the incredible capacity to rapidly alkalinize its external environment via amino acid catabolism (21–23). This process is primarily driven during carbohydrate stress by sensing of amino acids in the milieu and regulating amino acid import, largely governed by transcription factor Stp2p. Ammonia is extruded from the cell as these peptide substrates are consumed by the fungus, ultimately raising the extracellular pH (24, 25). Therefore, we utilized an *stp2Δ/Δ* mutant and isogenic revertant strain (*stp2Δ/Δ*+*STP2*) to determine the impact of alkalinization during coculture with *S. aureus*. The culture pH of the *stp2Δ/Δ* mutant during coculture was significantly decreased compared to that of the wild-type or revertant strains (Fig. 4A). In a similar fashion, production of alpha-toxin was significantly attenuated during coculture with the *stp2Δ/Δ* strain (Fig. 4B). Thus,




FIG 4 The alkalization-deficient *C. albicans* *stp2* Δ / Δ mutant fails to enhance *S. aureus* toxin production. Polymicrobial cultures of *S. aureus* and wild-type *C. albicans* (CA+SA), SA and *stp2* mutant (*stp2* Δ / Δ +SA), or SA and *STP2* revertant (*stp2* Δ / Δ +*STP2*+SA) were incubated at 37°C in media with an initial pH of 6. At 24 h, the pH of the cultures were measured using a pH meter (A), and alpha-toxin levels in supernatants were determined by ELISA (B). (C) Aliquots from each culture were taken at the endpoint to measure microbial counts by plating on selective media. Blue bars indicate *C. albicans* counts, and yellow bars indicate *S. aureus* counts. Solid color bars depict counts from monomicrobial (mono) cultures, while hatched bars depict counts from polymicrobial (poly) cultures. Data are representative of three independent repeats and expressed as the means plus SEM. Data were assessed for significance using one-way ANOVA and Dunnet’s posttest. *, $P < 0.05$; **, $P < 0.01$.

active alkalization of the external environment by *C. albicans* partially modulates staphylococcal alpha-toxin production during coculture.

Candida species differentially modulate alpha-toxin production during coculture. We next questioned whether augmented alpha-toxin production was specific to *C. albicans* or whether other non-*albicans* *Candida* (NAC) species could also potentiate this effect. Therefore, *S. aureus* was cultivated by itself or in the presence of various *Candida* species, including *C. albicans*, *C. glabrata*, *C. dubliniensis*, *C. tropicalis*, *C. parapsilosis*, and *C. krusei*. Measurement of the extracellular pH following coculture with *C. albicans*, *C. tropicalis*, and *C. krusei* revealed significantly increased neutralization compared to *S. aureus* monoculture (Fig. 5A). Coculture with *C. glabrata*, *C. dubliniensis*, and *C. parapsilosis* demonstrated only modest increases in pH which were generally below the threshold for robust *agr* activation. These results were recapitulated by examining alpha-toxin production during coculture. *Candida* species capable of significantly raising the extracellular pH in this assay (*C. albicans*, *C. tropicalis*, and *C. krusei*) also demonstrated increased capacity to augment alpha-toxin release, while those




FIG 5 *Candida* spp. differ in their capacity to modulate extracellular pH and augment toxin production during polymicrobial growth with *S. aureus*. Monomicrobial or polymicrobial cultures of *Candida* species (*C. albicans* [CA], *C. glabrata* [CG], *C. dubliniensis* [CD], *C. tropicalis* [CT], *C. parapsilosis* [CP], and *C. krusei* [CK]) and *S. aureus* (SA) were incubated at 37°C. After 16 h of growth, the pH of the cultures was measured using a standard pH meter (A), and the level of alpha-toxin in the supernatant was assessed by ELISA (B). Data are representative of five independent repeats and expressed as the means plus SEM. Data were assessed for significance using one-way ANOVA and Dunnet’s posttest. **, $P < 0.01$; ***, $P < 0.001$.

incapable of significantly modulating the extracellular pH demonstrated only modest elevation of this virulence determinant (Fig. 5B). These results demonstrate that exacerbation of alpha-toxin production is not limited to *C. albicans*, but this *in vitro* phenotype is largely driven by modulation of extracellular pH during coculture.

DISCUSSION

Often, the pathogenic process is delineated as the interaction between host and microbe resulting in some level of host detriment. Unfortunately, the contribution of microenvironment is all too frequently disregarded in this description. However, changes in environmental factors can ultimately drive or control pathogenicity or host susceptibility. Through the prism of single microbe infections, this concept seems obvious. For example, gene expression exhibited by *C. albicans* is very different in the gut compared to in the oral cavity—largely attributable to variations in host cell type, oxygen tension, nutrient availability, and microbial competition (26–28). In the context of a polymicrobial model of disease, gene regulation in response to environment by one organism may elicit reactions by a second pathogen, opportunist, or colonizer. An excellent example of this is the capacity of the lactobacilli to ferment carbohydrates to lactic acid at the vaginal mucosa (29). Production of relatively high levels of lactic acid keeps the vaginal pH low and consequently impairs the overgrowth of *C. albicans*, limiting the development of vulvovaginal candidiasis. While there are several examples of “environmental cross talk” resulting in microbial antagonism, reports regarding exacerbated virulence are less common (30–33).

We, along with others, have previously shown that murine intra-abdominal coinfection with *C. albicans* and *S. aureus* results in a strikingly high mortality rate (~90%) within ~16 h postinfection (p.i.), while infection with either microbe alone is nonlethal (17). Moreover, this synergistic lethality is dependent on alpha-toxin, as an isogenic strain lacking the gene encoding this virulence factor (*hla*) or neutralization of this toxin using a high-affinity antibody resulted in significant protection. In support of this, lavage of the peritoneal cavity following infection revealed ~4-fold-higher levels of alpha-toxin in coinfecting mice compared to mice challenged with *S. aureus* alone (17). These *in vivo* results substantially mimic those observed in the *in vitro* system used in this study.

Using several functional assays, we confirmed that alpha-toxin levels are significantly elevated during polymicrobial growth compared to monomicrobial growth in an *agr*-dependent fashion. Interestingly, the mechanism of alpha-toxin enhancement was not likely due to direct fungal ligand binding or recognition by the *agr* quorum sensing system, as *S. aureus* mutants with disrupted *agr* genes ($\Delta agrA$ or $\Delta agrB$) failed to demonstrate toxin production even during coculture with *C. albicans*, indicating the necessity of an intact *agr* regulon (Fig. 2). There are numerous examples where secreted microbial peptides or metabolites can exhibit a diverse array of effects on neighboring cells of the same or different species (30, 33, 34). Deletion of *agrB*, the autoinducing peptide permease, should still allow for functional *agr* signaling in the context of candidal factors capable of inducing the AgrC-AgrA two-component system. The observed failure to activate toxigenic and *agr* responses under such conditions or with spent *C. albicans* culture supernatant likely indicates that potential fungal ligands do not directly signal via the AgrC surface receptor.

The exclusion of a factor directly engaging *agr* led us to investigate potential indirect influences, such as modulation of environmental factors. In pioneering work characterizing the *agr* quorum sensing system in *S. aureus*, Regassa et al. demonstrated the pH responsiveness of the *agr* system. They reported increased *agr* activity during growth at pH 6.5 to 7 compared to growth in more alkaline or acidic media (15, 16). The pH of monomicrobial *S. aureus* cultures demonstrated significant acidification (pH 5.0 to 5.5), while cocultures exhibited a pH of 6.5 to 7, exactly in the range of maximum *agr* activity (Fig. 3A). Buffering of the media demonstrated that *agr* signaling could be manipulated positively or negatively during coculture depending on pH selection, further demonstrating the importance of environmental factors (e.g., pH) in driving

pathogenicity mechanisms *in vitro*. Both *C. albicans* and *S. aureus* preferentially utilize glucose to undergo oxidative and fermentative metabolism, ultimately producing acidic end products that drive lower culture pH. So then why does the culture medium containing *C. albicans* demonstrate an elevated pH? The ability of *C. albicans* (and other fungal species) to alkalinize its environment has been well documented in the literature. Vylkova and Lorenz have demonstrated that *C. albicans* can raise the pH of macrophage phagosomes, allowing for hyphal growth that damages the phagosomal membrane, aiding in escape and continued proliferation (21, 22). Moreover, the acidic pH (≤ 5) of the phagosome is vital for the activity of degradative enzymes that act to kill and digest engulfed pathogens, including fungi. Additionally, low pH is known to repress the morphological transition of yeast to hyphae, the major virulence attribute of *C. albicans* (35). Therefore, *C. albicans* has likely evolved strategies to modulate the external pH to bypass host checkpoints and killing mechanisms.

We propose that during polymicrobial growth, the pH initially decreases as glucose is metabolized by both organisms, then when glucose is limiting, the culture medium is alkalinized by *C. albicans*, in turn activating the *agr* quorum sensing system and subsequent alpha-toxin production. The mechanism of alkalinization involves the breakdown of amino acids for a carbon source and the subsequent excretion of ammonia, which raises the pH. In glucose-limited environments, amino acids are sensed by the SPS sensor system, a complex of three proteins (Ssy1, Ptr3, and Ssy5). This sensor complex induces the proteolytic cleavage of a cytoplasmic retention signal of Stp2p, allowing it to translocate to the nucleus. Stp2p binds SPS sensor-regulated promoters of various amino acid permease genes, including *CAN1*, *GAP1*, and *GAP2*, which transport extracellular amino acids into the cell (36). Amino acid catabolism begins with the deamination of an amino acid, catalyzed by amino acid-specific deaminases. The carbon backbone is converted to tricarboxylic acid (TCA) cycle intermediates (pyruvate, α -ketoglutarate, and acetoacetyl coenzyme A [acetoacetyl-CoA]) via the production of acetyl-CoA. The nitrogen is converted to ammonia and CO₂ by the urea amidolyase Dur1,2p, and subsequently excreted from the cell through various Ato (ammonia transport outward) family transmembrane proteins (23, 25, 36). *C. albicans* strains lacking *STP2* have an impaired capacity to alkalinize the environment compared to isogenic controls, corresponding with a decrease in ammonia produced during growth. Additionally, this mutant is unable to form hyphae after phagocytosis, preventing escape from the phagosome, which also leads to more effective killing by the macrophage. In a mouse model of disseminated candidiasis, *stp2 Δ / Δ* mutants displayed attenuated virulence as mortality was significantly delayed compared to wild-type or complemented strains (22). Using a *stp2 Δ / Δ* strain in our *in vitro* polymicrobial culture system, we found that the mutant cannot raise the pH to levels observed with the wild-type or revertant strains. This further confirms that the capacity of *C. albicans* to alkalinize the media contributes to enhanced alpha-toxin production during coculture (Fig. 4).

Although Stp2p is largely responsible for driving the alkalinization phenotype in *C. albicans*, it is not the sole mechanism for pH modulation. A number of *C. albicans* genes have been identified as having effects on external alkalinization, including *ALI1*, *SIN3*, *COX4*, *PEP8*, *KIS1*, and *CPH1*. Some of these genes (*COX4* and *KIS1*) are linked to carbon metabolism, while *CPH1* can regulate galactose utilization. *C. albicans* has also been shown to modulate pH without the production and excretion of ammonia, as seen during growth with nonfermentable carbon sources, like the carboxylic acids α -ketoglutarate, pyruvate, and lactate (24, 37). Interestingly, alkalinization occurs in low-glucose environments, regardless of the mechanism, indicating that this effect is glucose repressible. Vylkova and Lorenz hypothesize that alkalinization of the phagosome is due to limited glucose within the phagosomal compartment (22). Although glucose or other metabolic compounds were not a direct focus of this paper, their potential effect on pH and alpha-toxin production can be inferred. In addition to being responsive to pH, the *agr* quorum sensing system is known to be regulated by glucose levels, with growth under high glucose conditions correlated to low *agr* activity. Future

studies focusing on the metabolic profiles of both *C. albicans* and *S. aureus* may shed more light on this potential effector mechanism.

Last, we investigated the ability of other NAC species to enhance alpha-toxin production during *in vitro* growth. We found that the species differ in their alkalinization potential, with only *C. tropicalis* and *C. krusei* able to raise the pH of the media to levels similar to that of *C. albicans* during polymicrobial growth with *S. aureus*. Additionally, these three species were the only ones that caused a significant increase in the amount of alpha-toxin produced compared to *S. aureus* monomicrobial culture (Fig. 5). These data correlate nicely with work performed by Nash et al. in which mice were coinfecting with the exact same *Candida* strains. Mortality was observed in mice infected with *S. aureus* and *C. albicans*, *C. tropicalis*, and *C. krusei*. Strikingly, the species unable to modulate the pH and alpha-toxin production during *in vitro* growth in this study (*C. glabrata*, *C. dubliniensis*, and *C. parapsilosis*) were also nonlethal in the polymicrobial IAI model (38). A retrospective analysis of intra-abdominal infections, specifically looking at candidiasis, revealed that *C. albicans* is the most commonly isolated species (50 to 75%). The next most commonly isolated species is *C. glabrata* (12 to 25%), followed by *C. parapsilosis* (3 to 10%), *C. tropicalis* (3 to 5%), and *C. krusei* (3 to 6%) (39, 40). Interestingly, *C. glabrata* was implicated as the infectious organism in the majority (64%) of recurrent or persistent infections (39). Cheng et al. described persistent *C. glabrata* IAI in mice and an association with the formation of abscesses (41).

Various niches within the body are maintained at drastically different pH values. For example, the oral cavity is maintained at pH 6.2 to 7.6 by saliva (42). Blood is very tightly buffered to remain at pH 7.4, and a decrease in 0.05 units causes severe physiological problems, as seen in diabetic ketoacidosis (43). The vagina is more acidic, with a healthy pH ranging from 3.8 to 4.5 (44). The fluid within the peritoneal cavity of humans is reported to be at a pH of 7.5 to 8 (45). Although widely variable, bodily pH is tightly regulated to maintain homeostasis, and dysregulation is often indicative of poor health. A study investigating the transcriptome of *C. albicans* during murine IAI found that a number of genes involved in pH response were among the most highly upregulated. These genes include *RIM101*, the alkaline pH-regulated transcription factor that modulates morphology and gene expression (26). Indeed, a *rim101Δ/Δ* strain exhibited attenuated virulence during peritoneal infection, characterized by significantly lower fungal burdens (26). These findings indicate that the murine peritoneal cavity is alkaline and that *C. albicans* requires adaptation to pH to establish pathogenicity in this biological niche. However, because of the homeostatic nature of pH maintenance, it is unlikely that *C. albicans* is able to drastically alter the global pH of the peritoneal cavity during IAI. Our attempts to experimentally monitor broad pH changes in the peritoneal lavage fluid during mono- or coinfection have demonstrated insignificant differences, partially due to sensitivity of the techniques utilized (e.g., phenol red lavage, micro pH electrode) or spatiotemporal kinetics (data not shown). Thus, it is more likely that microenvironmental pH regulation, in peritoneal abscesses or tissues, may play an important role in driving these phenotypes. As *C. albicans* and *S. aureus* are known to tightly associate via staphylococcal binding of fungal hyphae via the candidal adhesin Als3p (46), it is possible that elevated *S. aureus* alpha-toxin production *in vivo* is due to local pH changes surrounding fungal-bacterial aggregates. However, this hypothesis requires future investigation.

Collectively, results from this study highlight the dynamic and complex nature of this fascinating microbial pair and polymicrobial interactions in general. Furthermore, they underscore the importance of environmental adaptation and its intersection with virulence that must be considered in the context of coinfection.

MATERIALS AND METHODS

Strains and growth conditions. *Candida albicans* SC5314 (referred to as CA) was used as the wild-type isolate throughout this work. The alkalinization-deficient mutant *stp2Δ/Δ* (SVC17) and its isogenic revertant strain *stp2Δ/Δ+STP2* (SVC19) were kind gifts from Michael Lorenz (University of Texas Health Science Center) (22). The non-*albicans* *Candida* (NAC) species *C. glabrata* CBS138 (CG), *C. dubliniensis* CD36 (CD), *C. parapsilosis* CDC317 (CP), *C. tropicalis* MYA3404 (CT), and *C. krusei* 81-B-5 (CK)

were used as species-representative strains (38). JE2, a USA300 isolate, was obtained from the Biodefense and Emerging Infectious (BEI) Research Resources repository and used as the wild-type *S. aureus* strain in this work and is referred to as SA. A *S. aureus* reporter strain [*S. aureus*(pDB22)] containing plasmid pDB22 (containing the P3 promoter fused to GFP_{mut2} and an erythromycin resistance cassette) was also used in this work (47). All strains were maintained as 20% glycerol stocks at -80°C .

Candida strains were streaked onto yeast-peptone-dextrose (YPD) agar and grown at 30°C . A single colony was inoculated in YPD broth and incubated overnight at 30°C with shaking at 200 rpm. *S. aureus* strains were streaked on Trypticase soy agar (TSA) (with $10\ \mu\text{g}/\text{ml}$ erythromycin added as needed). Single colonies were inoculated in Trypticase soy broth (TSB) and grown overnight at 37°C with shaking at 200 rpm. Aliquots ($500\ \mu\text{l}$) were washed three times with phosphate-buffered saline (PBS), and cell concentrations were adjusted to 1×10^7 CFU/ml. A 1:100 dilution was made into 5 ml of $0.6\times$ TSB plus 0.2% glucose (TSB-g) with the following groups: CA or SA (monomicrobial) and CA plus SA (CA+SA) (polymicrobial). PBS ($50\ \mu\text{l}$) was added to monomicrobial cultures. Cultures were incubated at 37°C with shaking at 200 rpm, and aliquots were removed at 16 h postinoculation (p.i.). Mono- and polymicrobial cultures using the alkalization-deficient mutant and revertant strains were prepared as described above, except TSB-g was adjusted to pH 6 prior to inoculation and cultures were incubated for 24 h.

Agr reporter assay. Monomicrobial and polymicrobial cultures were prepared as described above using CA and/or *S. aureus*(pDB22) ($10\ \mu\text{g}/\text{ml}$ erythromycin added for plasmid maintenance). At 16 h p.i., $100\text{-}\mu\text{l}$ aliquots were removed in triplicate from cultures and added to wells of 96-well black microtiter plate. Fluorescence (488 nm excitation, 525 nm emission) was measured using a plate reader (Synergy, Biotek). Experiments were repeated in triplicate, and results are expressed as the mean arbitrary fluorescence units (AFU) \pm standard errors of the means (SEM).

Blood agar lysis assay. Cultures were prepared as described above, and at 16 h p.i., 5-ml aliquots were centrifuged at 5,000 rpm to pellet cells, and supernatants were sterilized using $0.2\text{-}\mu\text{m}$ syringe filters. Sterile supernatants were concentrated $20\times$ by ethanol precipitation. Holes were punched in blood agar plates (TSA with 5% sheep blood) using a sterile pipette tip. Concentrated supernatant ($20\ \mu\text{l}$) was added to wells, and plates were incubated at 37°C for 24 h. Plates were photographed using a digital scanner (EPSON Perfection V700 Photo).

pH buffering. pH buffering experiments were done by the same culture set-up described above, with the following changes: 100 mM morpholinepropanesulfonic acid (MOPS) was added to $0.6\times$ TSB plus 0.2% glucose, and the medium was adjusted to various pHs (5.5, 6, 6.5, 7, 7.5, and 8) using 5 N HCl or NaOH as required. Aliquots ($500\ \mu\text{l}$) of overnight cultures of CA and *S. aureus*(pDB22) were washed with PBS and adjusted to 1×10^7 CFU/ml. A 1:100 dilution was made into 5 ml MOPS-buffered TSBg at each pH tested. Monomicrobial and polymicrobial cultures were incubated for 16 h at 37°C with shaking at 200 rpm.

Kinetic agr activation assay. *S. aureus*(pDB22) was grown overnight in MOPS-buffered $0.6\times$ TSB-g (pH 5.5) to inactivate *agr*. Additionally, *S. aureus* (JE2) and *C. albicans* were grown overnight in MOPS-buffered $0.6\times$ TSB-g (pH 7). Cultures were centrifuged at 4,000 rpm for 3 min. Supernatant from the *S. aureus* JE2 and *C. albicans* cultures were collected and filter sterilized. Cells from *S. aureus*(pDB22) grown at pH 5.5 were washed with PBS and adjusted to 1×10^{10} CFU/ml. Cells were added to 1 ml of $5\times$ MOPS-buffered TSB-g (pH 7) to a final concentration of 2×10^8 CFU/ml. Four milliliters of either *S. aureus* (JE2) spent medium, *C. albicans* spent medium, or fresh medium was added to the cells along with $10\ \mu\text{g}/\text{ml}$ erythromycin (for plasmid maintenance). Cultures were incubated at 37°C with shaking at 200 rpm. At 20-min intervals, $100\text{-}\mu\text{l}$ aliquots were removed, and fluorescence was measured as described above. Experiments were repeated in triplicate, and results are expressed as the mean AFU \pm SEM.

Alpha-toxin ELISA. *S. aureus* alpha-toxin was quantified by an enzyme-linked immunosorbent assay (ELISA), as described previously (17). Briefly, 96-well plates were coated with $0.1\ \mu\text{g}/\text{ml}$ MEDI4893* diluted in coating buffer and incubated overnight at 4°C . All wash steps were carried out with PBS containing 0.05% Tween 20. The plates were washed and blocked with SuperBlock (Pierce) for 1 h at room temperature. Fifty microliters of diluted filter-sterilized supernatant (taken at 16 h p.i.) was added to wells, with serial dilutions of native alpha-toxin included as the standard curve. The plates were incubated for 1 h at room temperature and then washed. Affinity-purified rabbit polyclonal anti-alpha-toxin antibody ($2\ \mu\text{g}/\text{ml}$) was added to wells, and the plates were incubated for 1 h at room temperature. The plates were washed, and then a 1:10,000 dilution of Affinipure horseradish peroxidase (HRP)-coupled goat anti-rabbit IgG detection antibody (Jackson Immuno Research) was added. The plates were incubated for 1 h at room temperature and washed. 3,3',5,5'-Tetramethylbenzidine (TMB) substrate was added to wells, and color was allowed to develop in the dark for 10 min. One hundred microliters of ELISA stop solution ($0.2\ \text{M}\ \text{H}_2\text{SO}_4$) was added to each well, and wells were read at 450 nm using a spectrophotometer (Synergy, Biotek). Culture supernatants from *S. aureus* NE1354 Δhla were used as background controls and were subtracted from sample wells to exclude any nonspecific binding of antibody by protein A. The experimental values were extrapolated to the standard curve. Experiments were completed in triplicate and shown as means \pm SEM.

CFU analysis. CFU enumeration was done by serial plating of culture media onto YPD containing $20\ \mu\text{g}/\text{ml}$ nafcillin (for *Candida* enumeration) and TSA containing $2.5\ \mu\text{g}/\text{ml}$ amphotericin B (for *S. aureus* enumeration) via the drop-plate method (48). The plates were incubated overnight at 37°C , and the bacteria were enumerated and expressed as CFU/milliliter. CFU values are representative of three independent repeats and represented as median \pm SEM.

ACKNOWLEDGMENTS

We thank Michael Lorenz (University of Texas Health Science Center) for providing strains SCV17 and SCV19. We also thank Pete Greenberg (University of Washington) for P3-GFP reporter plasmid pDB22. We gratefully acknowledge Biodefense and Emerging Infections Research Resources Repository (BEI Resources) for providing staphylococcal strains used in this work. Last, we thank Bret Sellman and Christine Tkaczyk (Astrazeneca) for providing MEDI-4893* and ELISA protocol.

This research was funded by National Institutes of Health, National Institute for Allergy and Infectious Disease, grants R01AI116025 (M.C.N., B.M.P.) and R01AI134796 (B.M.P.).

O.A.T. performed the experiments. B.M.P. and M.C.N. conceptualized the experiments. All authors contributed to writing, reviewing, and editing the manuscript.

We declare that we have no conflicts of interest.

The funders had no role in the design of the study, in the collection, analyses, or interpretation of data, in the writing of the manuscript, or in the decision to publish the results.

REFERENCES

- Carey AJ, Saiman L, Polin RA. 2008. Hospital-acquired infections in the NICU: epidemiology for the new millennium. *Clin Perinatol* 35:223–249. <https://doi.org/10.1016/j.clp.2007.11.014>.
- Klotz SA, Chasin BS, Powell B, Gaur NK, Lipke PN. 2007. Polymicrobial bloodstream infections involving *Candida* species: analysis of patients and review of the literature. *Diagn Microbiol Infect Dis* 59:401–406. <https://doi.org/10.1016/j.diagmicrobio.2007.07.001>.
- Valenza G, Tappe D, Turnwald D, Frosch M, König C, Hebestreit H, Abele-Horn M. 2008. Prevalence and antimicrobial susceptibility of microorganisms isolated from sputa of patients with cystic fibrosis. *J Cyst Fibros* 7:123–127. <https://doi.org/10.1016/j.jcf.2007.06.006>.
- Pate JC, Jones DB, Wilhelmus KR. 2006. Prevalence and spectrum of bacterial co-infection during fungal keratitis. *Br J Ophthalmol* 90: 289–292. <https://doi.org/10.1136/bjo.2005.081869>.
- Tawara Y, Honma K, Naito Y. 1996. Methicillin-resistant *Staphylococcus aureus* and *Candida albicans* on denture surfaces. *Bull Tokyo Dent Coll* 37:119–128.
- Gupta N, Haque A, Mukhopadhyay G, Narayan R, Prasad RK. 2005. Interactions between bacteria and *Candida* in the burn wound. *Burns* 31:375. <https://doi.org/10.1016/j.burns.2004.11.012>.
- Lopez N, Kobayashi L, Coimbra R. 2011. A comprehensive review of abdominal infections. *World J Emerg Surg* 6:7. <https://doi.org/10.1186/1749-7922-6-7>.
- de Ruiter J, Weel J, Manusama E, Kingma WP, van der Voort PH. 2009. The epidemiology of intra-abdominal flora in critically ill patients with secondary and tertiary abdominal sepsis. *Infection* 37:522–527. <https://doi.org/10.1007/s15010-009-8249-6>.
- McNeil JC, Vallejo JG, Hulten KG, Kaplan SL. 2018. Osteoarticular infections following open or penetrating trauma in children in the post-community-acquired methicillin-resistant *Staphylococcus aureus* era: the impact of *Enterobacter cloacae*. *Pediatr Infect Dis J* 37:1204–1210. <https://doi.org/10.1097/INF.0000000000001991>.
- Carlson E. 1982. Synergistic effect of *Candida albicans* and *Staphylococcus aureus* on mouse mortality. *Infect Immun* 38:921–924.
- Carlson E. 1983. Enhancement by *Candida albicans* of *Staphylococcus aureus*, *Serratia marcescens*, and *Streptococcus faecalis* in the establishment of infection in mice. *Infect Immun* 39:193–197.
- Carlson E. 1983. Effect of strain of *Staphylococcus aureus* on synergism with *Candida albicans* resulting in mouse mortality and morbidity. *Infect Immun* 42:285–292.
- Le KY, Otto M. 2015. Quorum-sensing regulation in staphylococci—an overview. *Front Microbiol* 6:1174. <https://doi.org/10.3389/fmicb.2015.01174>.
- Janzon L, Arvidson S. 1990. The role of the delta-lysin gene (*hld*) in the regulation of virulence genes by the accessory gene regulator (*agr*) in *Staphylococcus aureus*. *EMBO J* 9:1391–1399. <https://doi.org/10.1002/j.1460-2075.1990.tb08254.x>.
- Regassa LB, Novick RP, Betley MJ. 1992. Glucose and nonmaintained pH decrease expression of the accessory gene regulator (*agr*) in *Staphylococcus aureus*. *Infect Immun* 60:3381–3388.
- Regassa LB, Betley MJ. 1992. Alkaline pH decreases expression of the accessory gene regulator (*agr*) in *Staphylococcus aureus*. *J Bacteriol* 174:5095–5100. <https://doi.org/10.1128/jb.174.15.5095-5100.1992>.
- Todd OA, Fidel JPL, Harro JM, Hilliard JJ, Tkaczyk C, Sellman BR, Noverr MC, Peters BM. 2019. *Candida albicans* augments *Staphylococcus aureus* virulence by engaging the staphylococcal *agr* quorum sensing system. *mBio* 10:e00910-19. <https://doi.org/10.1128/mBio.00910-19>.
- Thoendel M, Kavanaugh JS, Flack CE, Horswill AR. 2011. Peptide signaling in the staphylococci. *Chem Rev* 111:117–151. <https://doi.org/10.1021/cr100370n>.
- Berube BJ, Bubeck-Wardenburg J. 2013. *Staphylococcus aureus* α -toxin: nearly a century of intrigue. *Toxins (Basel)* 5:1140–1166. <https://doi.org/10.3390/toxins5061140>.
- Somerville GA, Proctor RA. 2009. At the crossroads of bacterial metabolism and virulence factor synthesis in staphylococci. *Microbiol Mol Biol Rev* 73:233–248. <https://doi.org/10.1128/MMBR.00005-09>.
- Vylkova S, Lorenz MC. 2017. Phagosomal neutralization by the fungal pathogen *Candida albicans* induces macrophage pyroptosis. *Infect Immun* 85:e00832-16. <https://doi.org/10.1128/IAI.00832-16>.
- Vylkova S, Lorenz MC. 2014. Modulation of phagosomal pH by *Candida albicans* promotes hyphal morphogenesis and requires Stp2p, a regulator of amino acid transport. *PLoS Pathog* 10:e1003995. <https://doi.org/10.1371/journal.ppat.1003995>.
- Vylkova S, Carman AJ, Danhof HA, Collette JR, Zhou H, Lorenz MC. 2011. The fungal pathogen *Candida albicans* autoinduces hyphal morphogenesis by raising extracellular pH. *mBio* 2:e00055-11. <https://doi.org/10.1128/mBio.00055-11>.
- Danhof HA, Vylkova S, Vesely EM, Ford AE, Gonzalez-Garay M, Lorenz MC. 2016. Robust extracellular pH modulation by *Candida albicans* during growth in carboxylic acids. *mBio* 7:e01646-16. <https://doi.org/10.1128/mBio.01646-16>.
- Danhof HA, Lorenz MC. 2015. The *Candida albicans* ATO gene family promotes neutralization of the macrophage phagolysosome. *Infect Immun* 83:4416–4426. <https://doi.org/10.1128/IAI.00984-15>.
- Cheng S, Clancy CJ, Xu W, Schneider F, Hao B, Mitchell AP, Nguyen MH. 2013. Profiling of *Candida albicans* gene expression during intra-abdominal candidiasis identifies biologic processes involved in pathogenesis. *J Infect Dis* 208:1529–1537. <https://doi.org/10.1093/infdis/jit335>.
- Brown AJ, Odds FC, Gow NA. 2007. Infection-related gene expression in *Candida albicans*. *Curr Opin Microbiol* 10:307–313. <https://doi.org/10.1016/j.mib.2007.04.001>.
- Ghannoum MA, Jurevic RJ, Mukherjee PK, Cui F, Sikaroodi M, Naqvi A, Gillevet PM. 2010. Characterization of the oral fungal microbiome (mycobiome) in healthy individuals. *PLoS Pathog* 6:e1000713. <https://doi.org/10.1371/journal.ppat.1000713>.

29. Collins EB, Hardt P. 1980. Inhibition of *Candida albicans* by *Lactobacillus acidophilus*. *J Dairy Sci* 63:830–832. [https://doi.org/10.3168/jds.S0022-0302\(80\)83013-X](https://doi.org/10.3168/jds.S0022-0302(80)83013-X).
30. Wellington S, Greenberg EP. 2019. Quorum sensing signal selectivity and the potential for interspecies cross talk. *mBio* 10:e00146-19. <https://doi.org/10.1128/mBio.00146-19>.
31. Sztukowska MN, Dutton LC, Delaney C, Ramsdale M, Ramage G, Jenkinson HF, Nobbs AH, Lamont RJ. 2018. Community development between *Porphyromonas gingivalis* and *Candida albicans* mediated by InlJ and Als3. *mBio* 9:e00202-18. <https://doi.org/10.1128/mBio.00202-18>.
32. Madhani HD. 2011. Quorum sensing in fungi: Q&A. *PLoS Pathog* 7:e1002301. <https://doi.org/10.1371/journal.ppat.1002301>.
33. Abisado RG, Benomar S, Klaus JR, Dandekar AA, Chandler JR. 2018. Bacterial quorum sensing and microbial community interactions. *mBio* 9:e02331-17. <https://doi.org/10.1128/mBio.02331-17>.
34. Canovas J, Baldry M, Bojer MS, Andersen PS, Gless BH, Grzeskowiak PK, Stegger M, Damborg P, Olsen CA, Ingmer H. 2016. Cross-talk between *Staphylococcus aureus* and other staphylococcal species via the agr quorum sensing system. *Front Microbiol* 7:1733. <https://doi.org/10.3389/fmicb.2016.01733>.
35. Mayer FL, Wilson D, Hube B. 2013. *Candida albicans* pathogenicity mechanisms. *Virulence* 4:119–128. <https://doi.org/10.4161/viru.22913>.
36. Miramon P, Lorenz MC. 2016. The SPS amino acid sensor mediates nutrient acquisition and immune evasion in *Candida albicans*. *Cell Microbiol* 18:1611–1624. <https://doi.org/10.1111/cmi.12600>.
37. Miramon P, Lorenz MC. 2017. A feast for *Candida*: metabolic plasticity confers an edge for virulence. *PLoS Pathog* 13:e1006144. <https://doi.org/10.1371/journal.ppat.1006144>.
38. Nash EE, Peters BM, Fidel PL, Noverr MC. 2016. Morphology-independent virulence of *Candida* species during polymicrobial intra-abdominal infections with *Staphylococcus aureus*. *Infect Immun* 84:90–98. <https://doi.org/10.1128/IAI.01059-15>.
39. Vergidis P, Clancy CJ, Shields RK, Park SY, Wildfeuer BN, Simmons RL, Nguyen MH. 2016. Intra-abdominal candidiasis: the importance of early source control and antifungal treatment. *PLoS One* 11:e0153247. <https://doi.org/10.1371/journal.pone.0153247>.
40. Azim A, Ahmed A, Baronia AK, Marak RSK, Muzzafar N. 2017. Intra-abdominal candidiasis. *EMJ Nephrol* 5:83–92.
41. Cheng S, Clancy CJ, Hartman DJ, Hao B, Nguyen MH. 2014. *Candida glabrata* intra-abdominal candidiasis is characterized by persistence within the peritoneal cavity and abscesses. *Infect Immun* 82:3015–3022. <https://doi.org/10.1128/IAI.00062-14>.
42. Baliga S, Muglikar S, Kale R. 2013. Salivary pH: a diagnostic biomarker. *J Indian Soc Periodontol* 17:461–465. <https://doi.org/10.4103/0972-124X.118317>.
43. Kellum JA. 2000. Determinants of blood pH in health and disease. *Crit Care* 4:6–14. <https://doi.org/10.1186/cc644>.
44. Godha K, Tucker KM, Biehl C, Archer DF, Mirkin S. 2018. Human vaginal pH and microbiota: an update. *Gynecol Endocrinol* 34:451–455. <https://doi.org/10.1080/09513590.2017.1407753>.
45. Aoi W, Marunaka Y. 2014. Importance of pH homeostasis in metabolic health and diseases: crucial role of membrane proton transport. *BioMed Res Int* 2014:598986. <https://doi.org/10.1155/2014/598986>.
46. Peters BM, Ovchinnikova ES, Krom BP, Schlecht LM, Zhou H, Hoyer LL, Busscher HJ, van der Mei HC, Jabra-Rizk MA, Shirtliff ME. 2012. *Staphylococcus aureus* adherence to *Candida albicans* hyphae is mediated by the hyphal adhesin Als3p. *Microbiology* 158:2975–2986. <https://doi.org/10.1099/mic.0.062109-0>.
47. Yarwood JM, Bartels DJ, Volper EM, Greenberg EP. 2004. Quorum sensing in *Staphylococcus aureus* biofilms. *J Bacteriol* 186:1838–1850. <https://doi.org/10.1128/jb.186.6.1838-1850.2004>.
48. Donegan K, Matyac C, Seidler R, Porteous A. 1991. Evaluation of methods for sampling, recovery, and enumeration of bacteria applied to the phylloplane. *Appl Environ Microbiol* 57:51–56.