



A Point Mutation in the Transcriptional Repressor PerR Results in a Constitutive Oxidative Stress Response in *Clostridioides difficile* 630 Δ erm

Daniel Troitzsch,^a Hao Zhang,^b Silvia Dittmann,^a Dorothee Düsterhöft,^{a*} Timon Alexander Möller,^a Annika-Marisa Michel,^{b,d} Lothar Jänsch,^c Katharina Riedel,^a José Manuel Borrero-de Acuña,^{b,d}  Dieter Jahn,^{b,d}  Susanne Sievers^a

^aDepartment of Microbial Physiology and Molecular Biology, Institute of Microbiology, University of Greifswald, Greifswald, Germany

^bInstitute of Microbiology, Technical University of Braunschweig, Braunschweig, Germany

^cHelmholtz Center for Infection Research, Braunschweig, Germany

^dBraunschweig Integrated Centre of Systems Biology (BRICS), Technical University of Braunschweig, Braunschweig, Germany

ABSTRACT The human pathogen *Clostridioides difficile* has evolved into the leading cause of nosocomial diarrhea. The bacterium is capable of spore formation, which even allows survival of antibiotic treatment. Although *C. difficile* features an anaerobic lifestyle, we determined a remarkably high oxygen tolerance of the laboratory reference strain 630 Δ erm. A mutation of a single nucleotide (single nucleotide polymorphism [SNP]) in the DNA sequence (A to G) of the gene encoding the regulatory protein PerR results in an amino acid substitution (Thr to Ala) in one of the helices of the helix-turn-helix DNA binding domain of this transcriptional repressor in *C. difficile* 630 Δ erm. PerR is a sensor protein for hydrogen peroxide and controls the expression of genes involved in the oxidative stress response. We show that PerR of *C. difficile* 630 Δ erm has lost its ability to bind the promoter region of PerR-controlled genes. This results in a constitutive derepression of genes encoding oxidative stress proteins such as a rubrerythrin (*rbr1*) whose mRNA abundance under anaerobic conditions was increased by a factor of about 7 compared to its parental strain *C. difficile* 630. Rubrerythrin repression in strain 630 Δ erm could be restored by the introduction of PerR from strain 630. The permanent oxidative stress response of *C. difficile* 630 Δ erm observed here should be considered in physiological and pathophysiological investigations based on this widely used model strain.

IMPORTANCE The intestinal pathogen *Clostridioides difficile* is one of the major challenges in medical facilities nowadays. In order to better combat the bacterium, detailed knowledge of its physiology is mandatory. *C. difficile* strain 630 Δ erm was generated in a laboratory from the patient-isolated strain *C. difficile* 630 and represents a reference strain for many researchers in the field, serving as the basis for the construction of insertional gene knockout mutants. In our work, we demonstrate that this strain is characterized by an uncontrolled oxidative stress response as a result of a single-base-pair substitution in the sequence of a transcriptional regulator. *C. difficile* researchers working with model strain 630 Δ erm should be aware of this permanent stress response.

KEYWORDS *Clostridioides difficile*, oxidative stress response, transcriptional repressor, PerR, single nucleotide polymorphism, SNP, DNA binding, constitutive expression, rubrerythrin

Clostridioides difficile is a Gram-positive, anaerobic, spore-forming pathogen causing primarily hospital-acquired but increasingly also community-acquired infections, which has turned the bacterium into one of the most problematic

Citation Troitzsch D, Zhang H, Dittmann S, Düsterhöft D, Möller TA, Michel A-M, Jänsch L, Riedel K, Borrero-de Acuña JM, Jahn D, Sievers S. 2021. A point mutation in the transcriptional repressor PerR results in a constitutive oxidative stress response in *Clostridioides difficile* 630 Δ erm. *mSphere* 6:e00091-21. <https://doi.org/10.1128/mSphere.00091-21>.

Editor Craig D. Ellermeier, University of Iowa

Copyright © 2021 Troitzsch et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Susanne Sievers, susanne.sievers@uni-greifswald.de.

* Present address: Dorothee Düsterhöft, Max Planck Institute for Evolutionary Anthropology, Department of Evolutionary Genetics, Leipzig, Germany.

Received 31 January 2021

Accepted 5 February 2021

Published 3 March 2021

pathogens in human health care nowadays. *C. difficile* infections (CDIs) are often associated with high relapse rates and usually treated by broad-spectrum antibiotic therapy. Clinical symptoms of CDI vary from light diarrhea to acute infections like pseudomembranous colitis (1).

Due to its anaerobic lifestyle, oxygen (O₂) and reactive O₂ species in the human intestine represent a challenge for *C. difficile*. Remarkably, a high tolerance to O₂ was recently reported for a sporulation-deficient mutant of *C. difficile* 630Δ*erm* (2). Strain 630Δ*erm* is an erythromycin-sensitive and laboratory-generated derivative of the original patient-isolated strain 630 and is commonly used by *C. difficile* researchers as a reference strain for the generation of gene knockout mutants (3). Although the oxidative stress response is vital for an intestinal pathogen infecting its host, knowledge of the molecular details of oxidative adaptation mechanisms in *C. difficile* is still limited. Specifically, two general stress signatures of O₂ challenge of *C. difficile* strains 630 and 630Δ*erm* have been reported (4, 5), and for *C. difficile* 630Δ*erm*, an involvement of the alternative sigma factor σ^B was noted (6, 7). However, no sensor protein for oxidative conditions or mechanistic details on the regulation of the oxidative stress response in this important human pathogen have been described so far. Here, we report (i) an involvement of the hydrogen peroxide (H₂O₂) sensor and regulatory protein PerR in the oxidative stress response of *C. difficile* and (ii) defective repression by PerR and, thus, constitutive expression of oxidative stress genes in *C. difficile* reference strain 630Δ*erm*.

High abundance of oxidative stress proteins under anaerobic conditions. In previous studies, we observed a high abundance of oxidative stress-related proteins in *C. difficile* 630Δ*erm* already under conditions devoid of any oxidizing agents (8) and no significant induction when the bacterium was shifted to microaerobic conditions (5). Several of the corresponding genes are located at one genetic locus comprising a rubrerythrin (*rbr1*), the transcriptional repressor PerR (*perR*), a desulfoferrodoxin (*rbo*), and a glutamate synthase with an N-terminal rubredoxin fold (*CD630_08280*). Rbr1 even represents the second most abundant protein after the S-layer protein SlpA (8). We inquired why these genes are highly expressed in the absence of any oxidative stress and focused on the repressor protein PerR, which regulates its own transcription and that of genes involved in oxidative stress and metal homeostasis as described for *Bacillus subtilis* and other Gram-positive bacteria (9, 10). PerR is a member of the ferric uptake regulator (Fur) family and senses H₂O₂ stress by metal-catalyzed histidine oxidation (11) (Fig. 1A). Due to the permanently high cellular concentration of proteins encoded in the *rbr1* operon, we hypothesized a constitutive expression of the operon possibly caused by a failure of PerR-mediated gene repression under anaerobic conditions.

Amino acid substitution in the DNA binding domain of PerR. It was reported previously that the laboratory-generated strain *C. difficile* 630Δ*erm* features several genome alterations compared to its parental strain *C. difficile* 630 (12, 13). We aligned *perR* sequences of *C. difficile* 630 and *C. difficile* 630Δ*erm* and found a single nucleotide polymorphism (SNP) (A to G) resulting in an amino acid conversion from threonine to alanine at position 41. An alignment of the PerR amino acid sequence with sequences from 11 other clinically relevant *C. difficile* strains revealed that the T41A substitution is unique to laboratory strain 630Δ*erm* (see Fig. S1 in the supplemental material). A comprehensive sequence alignment of over 900 proteins of the Fur family from different species showed that the threonine at position 41 is highly conserved and present in over 80% of the investigated proteins. More than 90% of the proteins contain a threonine or serine at this position (Data Set S1). A structural comparison of previously investigated DNA binding domains of Fur and PerR homologues in *Escherichia coli*, *B. subtilis*, *Staphylococcus epidermidis*, and *Streptococcus pyogenes* to the *C. difficile* 630Δ*erm* PerR sequence indicates that the T41A mutation is located in a helix of the helix-turn-helix motif of the DNA binding domain (Fig. 1B and C). The DNA promoter sequences upstream of *rbr1* are identical between strains 630 and 630Δ*erm* (Fig. S2). We therefore hypothesized that the amino acid substitution in PerR is the reason for the loss of its binding to PerR boxes on the DNA and possibly causes increased O₂




FIG 1 DNA binding of PerR. (A) Mode of PerR function shown schematically. PerR bound to the DNA represses the expression of the *rbr1* operon. H_2O_2 treatment leads to PerR oxidation, a conformational change, and the release of the DNA promoter, resulting in the expression of the *rbr1* operon. (B) Structure of *C. difficile* 630Δerm PerR deduced from the homologue of *B. subtilis*. The DNA binding site, including the helix-turn-helix (HTH) motif and the dimerization domain, is marked. The location of the amino acid exchange is marked by a red box. (C) Amino acid sequence of the DNA binding site of *C. difficile* 630Δerm PerR. The T41A mutation is marked in red.

tolerance of strain 630Δerm compared to other *C. difficile* strains, including its parental strain 630.

Increased O_2 tolerance, derepression of the *rbr1* operon, and missing DNA binding of PerR. To investigate differences in O_2 tolerance between *C. difficile* 630 and 630Δerm, we counted CFU for both strains after cells had been exposed to atmospheric O_2 concentrations (Fig. 2A and Fig. S3). Strain 630 showed a significantly higher susceptibility to O_2 than its derivative 630Δerm, of which a substantial number of cells survived even after 9 h of challenge.

Furthermore, we were able to show an increased O_2 tolerance of *C. difficile* 630 after a 10-min adaptation treatment with 0.4 mM H_2O_2 or 5% O_2 . This experiment demonstrates that O_2 tolerance is indeed inducible in strain 630, possibly mediated by the inactivation of the repressor PerR (Fig. 2A). The direct oxidizing agent of the histidine residues in the DNA binding site of PerR was reported to be H_2O_2 . However, pretreatment with molecular O_2 also resulted in an increased O_2 tolerance of strain 630. Probably, molecular O_2 is converted to superoxide, which is subsequently reduced to H_2O_2 by a superoxide reductase such as Rbo. Noteworthy, adaptation with 0.4 mM H_2O_2 is quite stringent for *C. difficile* since absolute cell numbers are much lower (Fig. S3), and although the positive effect of adaptation with H_2O_2 is outstanding at 1 h and 3 h of O_2 challenge, it lasts for only a limited time, with almost no viable cells counted after 6 h and 9 h.

To prove that the higher O_2 tolerance of *C. difficile* 630Δerm was caused by the missing binding of PerR to its cognate DNA binding site, we analyzed transcriptional levels of the *rbr1* operon under H_2O_2 stress and control conditions (Fig. 2B and C and Fig. S4). Transcription of the operon was inducible by H_2O_2 in strain 630 by a factor of 2.4, whereas transcript levels in 630Δerm were permanently very high and not further inducible by H_2O_2 (factor of 0.96).

To verify that the increased O_2 tolerance of strain 630Δerm and its constitutive expression of the *rbr1* operon are the results of an inactive PerR protein, we constructed a *C. difficile* 630Δerm strain that is complemented with intact PerR of strain *C. difficile* 630 in *trans* and inducible by tetracycline (*C. difficile* DT022). Subsequently, we analyzed the PerR-repressed *rbr1* transcription of this strain (Fig. 2C). After induction with anhydrotetracycline (ATc), *rbr1* mRNA levels were reduced even below the ones of strain 630 (factor of 0.84), indicating that *in vivo* PerR activity and, consequently,




FIG 2 CFU counting, transcriptional analyses, and DNA-PerR interaction. (A) Survival in the presence of O₂ of *C. difficile* strains 630, 630 Δerm , and 630 adapted to 5% O₂ for 10 min or 0.4 mM H₂O₂ for 10 min. Ratios of aerobically cultured cells/cells of the corresponding cultures kept anaerobically are given for 1, 3, 6, and 9 h of O₂ challenge. * indicates statistical significance. (B) Northern blot analysis of the expression of the *rbr1* operon. Lane 1, 630 control; lane 2, 630 induced with H₂O₂; lane 3, 630 Δerm control; lane 4, 630 Δerm induced with H₂O₂. Total RNA levels were monitored by methylene blue (MB) staining. nt, nucleotide. (C) Transcription of the *rbr1* gene was quantified for *C. difficile* 630, 630 Δerm , and DT022 containing the plasmid pDSW1728-*perR* by RT-qPCR analysis and related to the *C. difficile* 630 control (CTL). ATc, induction with anhydrotetracycline. *, $P < 0.05$; ***, $P < 0.001$. (D) EMSAs were carried out with 2 ng of the *rbr1* promoter fragment (220 bp), concentrated at 147 nM, from strain *C. difficile* 630. In lanes 1 and 5, the DNA fragment was incubated without protein. The DNA fragment was incubated with purified PerR 630 (lanes 2 to 4) or PerR 630 Δerm (lanes 6 to 8) with increasing protein amounts (360 ng, 420 ng, and 480 ng, respectively).

DNA binding of PerR in *C. difficile* 630 Δerm were restored. Transcription in the complementation strain DT022 was inducible by H₂O₂ (factor of 3.52); i.e., the plasmid-encoded PerR of strain 630 responded to H₂O₂.

To confirm that the T41A exchange in PerR of strain 630 Δerm is the sole reason that hampers PerR box binding, we performed electrophoretic mobility shift assays (EMSAs) excluding any other cellular factors. PerR proteins from strains 630 and 630 Δerm were recombinantly produced, purified to apparent homogeneity (Fig. S5), and incubated with a labeled 220-bp upstream promoter fragment of *rbr1*. While PerR from strain 630 led to a clear shift of the DNA band, no shift was detectable for PerR from *C. difficile* 630 Δerm at any tested protein concentration (Fig. 2D).

Conclusion. This study demonstrated the constitutive derepression of genes involved in the oxidative stress response in strain *C. difficile* 630 Δerm caused by only

one SNP in the DNA sequence of the transcriptional repressor PerR. Researchers undertaking physiological studies in this strain should be aware of the permanent stress response mediated by missing PerR repression. Since the oxidative stress response is linked to other cellular networks, e.g., metal homeostasis and O₂-sensitive metabolic pathways, experiments carried out in strain 630 Δ erm might come to a different conclusion than experiments that would have been undertaken in any other *C. difficile* strain. Resistance to oxidative stress is vital for a pathogen to survive in the host. Hence, infection studies using the laboratory-constructed strain 630 Δ erm could also significantly vary from studies based on the original and patient-isolated strain. Since *C. difficile* 630 Δ erm is used as a reference strain in many laboratories for the construction of gene inactivation mutants, researchers should consider the background oxidative stress response when interpreting their experimental data.

Methods. (i) Bioinformatic methods. Structural analysis of the *C. difficile* PerR protein was performed using the Phyre2 Web portal (14). For the alignment of over 900 Fur family proteins, MView was used (15). The sequence alignments were carried out with Clustal Omega (16). PerR boxes were identified using Virtual Footprint Version 3.0 (17).

(ii) Plasmid and strain construction. *C. difficile* strains 630 and 630 Δ erm were obtained from the German Collection of Microorganisms and Cell Cultures GmbH (DSMZ) (Braunschweig, Germany). The *perR* gene was amplified from genomic DNA of *C. difficile* 630 using forward primer 5'-ATGGTAGAGCTCAATATAATGTTGGGAGGAATTAAAGAAATGAAATTTCTAAACAACGAG-3' and reverse primer 5'-ATGGTAGGATCCTTATTTCTCGAACTGCGGGTG-3' (Thermo Fisher Scientific, Waltham, MA, USA). The forward primer includes the ribosome binding site (RBS) of *C. difficile* SlpA and a SacI restriction site, and the reverse primer includes a BamHI restriction site. The PCR product was digested with SacI and BamHI (Thermo Fisher Scientific, Waltham, MA, USA) and ligated into the tetracycline-inducible vector pDSW1728 (18). pDSW1728 was a gift from Craig Ellermeier and David S. Weiss (Addgene plasmid 120812 [<http://n2t.net/addgene:120812>]; RRID, Addgene_120812). The resulting plasmid, pDSW1728-*perR*, was transformed into *E. coli* ST18 (19) following mating in *C. difficile* 630 Δ erm to construct *C. difficile* DT022. Sequences were checked by sequencing (see Fig. S6 in the supplemental material).

(iii) Bacterial strains and growth conditions. All *C. difficile* strains were cultured in brain heart infusion (BHI) medium as previously described (20), supplemented as needed with 25 μ g/ml thiamphenicol. For a first preculture, 200 μ l of spores was germinated in BHI medium supplemented with 0.1% cysteine. After 72 h, 10 ml of BHI medium was inoculated with 10 μ l of the first preculture to obtain a second preculture. This second preculture was directly diluted 1:10 in four steps and incubated overnight. The main cultures were inoculated to an A₆₀₀ of 0.05 from a second preculture exhibiting an A₆₀₀ of between 0.8 and 1.0. The expression of plasmid pDSW1728-*perR* was induced by adding 200 ng/ml anhydrotetracycline (ATc) at an A₆₀₀ of 0.1. For Northern blot and reverse transcription-quantitative PCR (RT-qPCR) analyses, *C. difficile* 630, *C. difficile* 630 Δ erm, and *C. difficile* DT022 were grown to the exponential growth phase to an A₆₀₀ of 0.4, before cultures were split and one of the two subcultures was stressed with 0.4 mM H₂O₂ for 10 min. Samples were taken to allow a later preparation of RNA (21). For CFU counting experiments, *C. difficile* 630 and 630 Δ erm were cultured to an A₆₀₀ of 0.4. The *C. difficile* 630 culture was split into three subcultures: one subculture was treated with 0.4 mM H₂O₂ for 10 min, the second subculture was flushed with 5% O₂, and the third subculture remained untreated (control). All cultures were centrifuged for 5 min at 8,500 rpm, and the supernatant was discarded. The cell pellets were resuspended in 30 ml fresh BHI medium. From each of the four cultures, 15 ml was transferred to two different 92- by 16-mm petri dishes. One petri dish was incubated aerobically, and the other was incubated anaerobically. Samples were taken before and 1, 3, 6, and 9 h after O₂ exposure in three biological replicates, and dilution series were plated on BHI agar plates and incubated anaerobically (Fig. S3). For the overexpression of PerR, *Escherichia coli* BL21 grown in LB medium was used.

(iv) RNA preparation. For cell lysis and RNA isolation, TRIzol reagent provided by Invitrogen (Thermo Fisher Scientific, Waltham, MA, USA) was used according to the manufacturer's protocol (22). RNA solubilized in diethyl pyrocarbonate (DEPC)-treated water was stored at -70°C .

(v) Transcriptional profiling. A PCR fragment of gene *rbr1* was prepared using chromosomal DNA of *C. difficile* 630 as a template with primers 5'-AATGGCAGGATTGTCAGGAG-3' and 5'-CTAATACGACTACTATAGGGAGATGGATGGTCACATACTGGGC-3'. Digoxigenin (DIG)-labeled RNA probes were obtained and Northern blot analyses were carried out as previously described (21). *rbr1* transcription was quantified by RT-qPCR in three biological replicates with three technical replicates each using the above-mentioned primers. The *rpoC* gene with forward primer 5'-CTAGCTGCTCCATATGCTCACATC-3' and reverse primer 5'-CCAGTCTCTCTGGATCAACTA-3' served as a reference. cDNA synthesis and qPCR were performed as described previously (23). The qPCRs were performed on a qTOWER 2.2 quantitative PCR thermocycler (Analytik Jena, Jena, Germany) (Fig. S4).

(vi) Statistical analyses. For CFU counting experiments, ratios of CFU counted under aerobic conditions versus anaerobic conditions were calculated. Differences between the ratios were tested for statistical significance using multiple *t* testing. α was set to 0.05 and was subsequently corrected using the Holm-Bonferroni method (24). The RT-qPCR quantitative data analysis was based on the Pfaffl method (25), and statistical analysis was performed using Student's *t* test.

Visualization of statistical analyses was performed using GraphPad Prism software (GraphPad Software Inc., La Jolla, CA).

(vii) Overexpression and purification of PerR of *C. difficile* 630 Δ erm in *E. coli* BL21. Protein overproduction was monitored as previously reported (26), with the following modifications. First, synthesized gene variants (Thermo Fisher Scientific, Waltham, MA, USA) were introduced into pGEX6P1 allowing for glutathione S-transferase (GST)-mediated affinity purification and subsequent tag excision. *E. coli* BL21 cells harboring recombinant genes were induced (0.1 mM isopropyl- β -D-thiogalactopyranoside [IPTG]) at an optical density at 600 nm (OD_{600}) of 0.5, grown aerobically for 4 h, and shifted to anaerobiosis for 2 h.

(viii) Electrophoretic mobility shift assay. Shift assays were conducted as previously specified (27), with minor variations. The 220-bp *rbr1* promoter region was amplified via PCR using forward primer 5'-TTGCAATAGGTATAGCGACAAG-3' and reverse primer 5'-TGCAATAGGTATAGCGACAAG-3'. The electrophoretic mobility shift assay (EMSA) was performed under anaerobic conditions.

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

FIG S1, PDF file, 0.2 MB.

FIG S2, PDF file, 0.4 MB.

FIG S3, PDF file, 0.3 MB.

FIG S4, PDF file, 0.2 MB.

FIG S5, PDF file, 0.1 MB.

FIG S6, PDF file, 0.1 MB.

DATA SET S1, PDF file, 5.5 MB.

ACKNOWLEDGMENTS

This work was funded by the German Research Foundation (231396381/GRK1947); the Ministerium für Bildung, Wissenschaft und Kultur Mecklenburg-Vorpommern (UG16001); and the Federal State of Lower Saxony, Niedersächsisches Vorab CDiff and CDInfect projects (VWZN2889/3215/3266). We acknowledge support for the article processing charge from the DFG (German Research Foundation) (393148499) and the Open Access Publication Fund of the University of Greifswald.

E. coli ST18 was a kind gift from the laboratory of J.M.B.-D.A.

REFERENCES

1. Smits WK, Lyras D, Lacy DB, Wilcox MH, Kuijper EJ. 2016. *Clostridium difficile* infection. Nat Rev Dis Primers 2:16020. <https://doi.org/10.1038/nrdp.2016.20>.
2. Edwards AN, Karim ST, Pascual RA, Jowhar LM, Anderson SE, McBride SM. 2016. Chemical and stress resistances of *Clostridium difficile* spores and vegetative cells. Front Microbiol 7:1698. <https://doi.org/10.3389/fmicb.2016.01698>.
3. Hussain HA, Roberts AP, Mullany P. 2005. Generation of an erythromycin-sensitive derivative of *Clostridium difficile* strain 630 (630Deltaerm) and demonstration that the conjugative transposon Tn916DeltaE enters the genome of this strain at multiple sites. J Med Microbiol 54:137–141. <https://doi.org/10.1099/jmm.0.45790-0>.
4. Emerson JE, Stabler RA, Wren BW, Fairweather NF. 2008. Microarray analysis of the transcriptional responses of *Clostridium difficile* to environmental and antibiotic stress. J Med Microbiol 57:757–764. <https://doi.org/10.1099/jmm.0.47657-0>.
5. Neumann-Schaal M, Metzendorf NG, Troitzsch D, Nuss AM, Hofmann JD, Beckstette M, Dersch P, Otto A, Sievers S. 2018. Tracking gene expression and oxidative damage of O₂-stressed *Clostridioides difficile* by a multi-omics approach. Anaerobe 53:94–107. <https://doi.org/10.1016/j.anaerobe.2018.05.018>.
6. Kint N, Janoir C, Monot M, Hoys S, Soutourina O, Dupuy B, Martin-Verstraete I. 2017. The alternative sigma factor σ B plays a crucial role in adaptive strategies of *Clostridium difficile* during gut infection. Environ Microbiol 19:1933–1958. <https://doi.org/10.1111/1462-2920.13696>.
7. Boekhoud IM, Michel A-M, Corver J, Jahn D, Smits WK. 2020. Redefining the *Clostridioides difficile* σ B regulon: σ B activates genes involved in detoxifying radicals that can result from the exposure to antimicrobials and hydrogen peroxide. mSphere 5:e00728-20. <https://doi.org/10.1128/mSphere.00728-20>.
8. Otto A, Maaß S, Lassek C, Becher D, Hecker M, Riedel K, Sievers S. 2016. The protein inventory of *Clostridium difficile* grown in complex and minimal medium. Proteomics Clin Appl 10:1068–1072. <https://doi.org/10.1002/prca.201600069>.
9. Dubbs JM, Mongkolsuk S. 2012. Peroxide-sensing transcriptional regulators in bacteria. J Bacteriol 194:5495–5503. <https://doi.org/10.1128/JB.00304-12>.
10. Hillmann F, Fischer R-J, Saint-Prix F, Girbal L, Bahl H. 2008. PerR acts as a switch for oxygen tolerance in the strict anaerobe *Clostridium acetobutylicum*. Mol Microbiol 68:848–860. <https://doi.org/10.1111/j.1365-2958.2008.06192.x>.
11. Lee J-W, Helmann JD. 2006. The PerR transcription factor senses H₂O₂ by metal-catalysed histidine oxidation. Nature 440:363–367. <https://doi.org/10.1038/nature04537>.
12. van Eijk E, Anvar SY, Browne HP, Leung WY, Frank J, Schmitz AM, Roberts AP, Smits WK. 2015. Complete genome sequence of the *Clostridium difficile* laboratory strain 630 Δ erm reveals differences from strain 630, including translocation of the mobile element CTn5. BMC Genomics 16:31. <https://doi.org/10.1186/s12864-015-1252-7>.
13. Coltery MM, Kuehne SA, McBride SM, Kelly ML, Monot M, Cockayne A, Dupuy B, Minton NP. 2017. What's a SNP between friends: the influence of single nucleotide polymorphisms on virulence and phenotypes of *Clostridium difficile* strain 630 and derivatives. Virulence 8:767–781. <https://doi.org/10.1080/21505594.2016.1237333>.
14. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 Web portal for protein modelling, prediction and analysis. Nat Protoc 10:845–858. <https://doi.org/10.1038/nprot.2015.053>.
15. Brown NP, Leroy C, Sander C. 1998. MView: a Web-compatible database search or multiple alignment viewer. Bioinformatics 14:380–381. <https://doi.org/10.1093/bioinformatics/14.4.380>.
16. Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN, Potter SC, Finn RD, Lopez R. 2019. The EMBL-EBI search and sequence analysis tools APIs in 2019. Nucleic Acids Res 47:W636–W641. <https://doi.org/10.1093/nar/gkz268>.
17. Münch R, Hiller K, Grote A, Scheer M, Klein J, Schobert M, Jahn D. 2005. Virtual Footprint and PRODORIC: an integrative framework for regulon prediction in prokaryotes. Bioinformatics 21:4187–4189. <https://doi.org/10.1093/bioinformatics/bti635>.
18. Ransom EM, Ellermeier CD, Weiss DS. 2015. Use of mCherry red fluorescent protein for studies of protein localization and gene expression in *Clostridium difficile*. Appl Environ Microbiol 81:1652–1660. <https://doi.org/10.1128/AEM.03446-14>.
19. Thoma S, Schobert M. 2009. An improved *Escherichia coli* donor strain for diparental mating. FEMS Microbiol Lett 294:127–132. <https://doi.org/10.1111/j.1574-6968.2009.01556.x>.
20. Sievers S, Dittmann S, Jordt T, Otto A, Hochgräfe F, Riedel K. 2018. Comprehensive redox profiling of the thiol proteome of *Clostridium difficile*. Mol Cell Proteomics 17:1035–1046. <https://doi.org/10.1074/mcp.TIR118.000671>.
21. Müller M, Reiß S, Schlüter R, Mäder U, Beyer A, Reiß W, Marles-Wright J, Lewis RJ, Pförtner H, Völker U, Riedel K, Hecker M, Engelmann S, Pané-Farré J. 2014. Deletion of membrane-associated Asp23 leads to upregulation of cell wall stress genes in *Staphylococcus aureus*. Mol Microbiol 93:1259–1268. <https://doi.org/10.1111/mmi.12733>.
22. Chomczynski P. 1993. A reagent for the single-step simultaneous isolation of RNA, DNA and proteins from cell and tissue samples. Biotechniques 15:532–534, 536–537.
23. Sievers S, Sternkopf Lillebæk EM, Jacobsen K, Lund A, Møllerup MS, Nielsen PK, Kallipolitis BH. 2014. A multicopy sRNA of *Listeria monocytogenes* regulates expression of the virulence adhesin LapB. Nucleic Acids Res 42:9383–9398. <https://doi.org/10.1093/nar/gku630>.
24. Holm S. 1979. A simple sequentially rejective multiple test procedure. Scand J Stat 6:65–70.
25. Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res 29:e45. <https://doi.org/10.1093/nar/29.9.e45>.
26. Hillmann F, Döring C, Riebe O, Ehrenreich A, Fischer R-J, Bahl H. 2009. The role of PerR in O₂-affected gene expression of *Clostridium acetobutylicum*. J Bacteriol 191:6082–6093. <https://doi.org/10.1128/JB.00351-09>.
27. Frädlich C, March A, Fiege K, Hartmann A, Jahn D, Härtig E. 2012. The transcription factor AlSR binds and regulates the promoter of the *alsSD* operon responsible for acetoin formation in *Bacillus subtilis*. J Bacteriol 194:1100–1112. <https://doi.org/10.1128/JB.06425-11>.